

## Pertuzumab and Trastuzumab

---

### What is Pertuzumab (per-TOO-zoo-mab) and how does it work?

Pertuzumab is a type of targeted cancer therapy called a “monoclonal antibody”. Another name for this medicine is Perjeta. Pertuzumab seeks out cancer cells and attaches to the HER2 receptors, to prevent the cells from dividing and making new cancer cells.

### What is Trastuzumab (tras-TOO-zoo-mab) and how does it work?

Trastuzumab is a type of targeted cancer therapy called a “monoclonal antibody”. Another name for this medicine is Herceptin. Trastuzumab seeks out cancer cells and attaches to the HER2 receptors, to prevent the cells from dividing and making new cancer cells. It also fights cancer cells by stimulating your immune system.

### Why am I getting 2 treatment medicines for my cancer?

Both medicines work to stop fast growing cancer cells from dividing and making new cells. Each medicine attacks the cancer cells differently. Giving the medicines together makes your treatment more effective.

### What should I tell my doctor before my treatment?

Talk to your doctor about the following:

- If you have ever had chemotherapy or anti-cancer treatment and the names of the medicines you were given.
- If you have ever had heart, liver, or lung problems.
- If you are allergic to mice.

---

**This handout is for informational purposes only. Talk with your doctor or health care team if you have any questions about your care.**

- If you are pregnant or think you may be pregnant. Your doctor will talk with you about birth control when you take these medicines.
- If you are breastfeeding.
- If you have been told that you need to start a new medicine.
- Other medicines/pills you take, including:
  - ▶ Medicines ordered by any of your doctors
  - ▶ Herbs
  - ▶ Vitamins
  - ▶ Over-the-counter medicines

## How does my doctor decide my treatment dose?

To determine your treatment dose, your doctor will review the following: your height, your weight, your medicines, how well your heart works and any other health problems you have.

You will receive a dose every 3 weeks. Normally, you will continue to receive this treatment as long as you do not have too many side effects and your cancer is not getting worse. You will see your doctor or nurse practitioner before each treatment.

## Should I eat or drink before my treatment?

It is best to eat a small meal and drink plenty of fluids before your treatment begins. If you have been told to limit fluids, check with your doctor about how much you can drink.

## How will my treatment be given?

**Pertuzumab** is given directly into your blood stream through a tube (IV) placed in your arm or chest. Pertuzumab comes in a bag with tubing attached. Your nurse will connect the tubing to a pump. Your first treatment takes 60 minutes. If you do not have any problems with your first treatment, then your future treatments will only take 30 minutes.

**Trastuzumab** is given directly into your blood stream through a tube (IV) placed in your arm or chest. Trastuzumab comes in a bag with tubing attached. Your nurse will connect the tubing to a pump. Your first treatment takes 90 minutes. If you do not have any problems with your first treatment, then your future treatments will only take 30 minutes.

Your entire visit, including your doctor's appointment and your treatment will take about 3 to 5 hours.

## What are the side effects of this treatment?

Every person responds differently to treatment. Some of the more common side effects of this treatment are:

- Diarrhea
- Fatigue
- Weakness
- Nausea
- Rash or other skin changes such as dryness or itching
- Flu-like symptoms: chills, fever, headache, body aches and sweating (these are more common with your first few doses of Trastuzumab)
- Treatment related reactions including: difficulty breathing, hives, or swelling of your face
- Cough

## When should I call my doctor?

You should call your doctor right away if you have any of the following signs or symptoms:

- Fever of 100.4 degrees Fahrenheit (38 degrees Celsius) or higher

**A fever can be life-threatening if not treated. Your doctor may ask you to go to the hospital.**

- Chills, sore throat, cough or a wound that does not get better
- Open sores in your mouth
- Nausea that prevents you from eating or drinking
- Vomiting even after you have taken your anti-nausea medicine
- Bleeding or bruising, including bloody or black stools or blood in your urine
- Diarrhea (4 or more loose stools in 24 hours)
- Changes in your heartbeat
- Shortness of breath or difficulty breathing
- Swelling in you feet or ankles
- Muscle or joint aches and pain
- Severe tiredness, weakness or pain

- **Signs of an allergic reaction, including:**
  - ▶ Difficulty breathing, wheezing, chest pain or tightness
  - ▶ Swelling in your face, lips, tongue, or throat
  - ▶ Rash

### **Is there anything else I should know about this treatment?**

- **Trastuzumab and Pertuzumab** may cause problems with the pumping of your heart. Your doctor will check your heart before you start these medicines and every several months while you are getting treatment.
- This treatment may change how your body reacts to vaccines. Talk to your doctor before getting any vaccines.
- It is important to keep hydrated during treatment. Unless told otherwise by your doctor, drink 8 to 10 cups of non-caffeinated fluid each day.

For more information about cancer, cancer treatment, side effects or how to care for yourself during treatment, ask your doctor, nurse or pharmacist.

You may also find it helpful to watch The James Patient Education videos at <http://cancer.osu.edu/patientedvideos> to help you learn tips for managing treatment side effects.